

**A remark to “A note on the Fundamental Group of a
Triangular Algebra”, by F. Xu.**

Juan Carlos Bustamante and Diane Castonguay

ABSTRACT. We provide a counter-example to Proposition 3.2 of [4].

In [1], the quiver Γ of homotopy relations of admissible presentations of an algebra (as quotients of some path algebra) was introduced. In [3], P. Le Meur showed that if $A \simeq kQ/I$ and

- I is a monomial ideal and
- the quiver Q has no multiple arrows

then the quiver Γ has a unique source, without any additional hypothesis about the characteristic of the field k .

Furthermore, in [2], P. Le Meur showed that if

- k is a field of characteristic zero, and
- the quiver Q has no double bypasses,

then the quiver Γ has a unique source.

These results ensure that under some hypotheses, there is a privileged homotopy relation, and consequently, a privileged fundamental group among all the fundamental groups that can arise as fundamental groups of some presentation of a given algebra.

In [2], Example 3, page 345, shows that the second quoted result is not true if one drops the two hypotheses simultaneously. In that example, which consists of a quiver having a double bypass, and considering a field of characteristic 2 one obtains a quiver Γ with two sources. Moreover, there is a suggested generalization of this example to any non-zero value of $\text{char } k$.

It is then natural to ask if one can drop one of the two hypotheses to generalize Le Meur’s result. The question has been tackled in [4]. In that paper, the framework is that of triangular algebras over fields of characteristic zero. One can find the following Proposition:

Proposition 3.2 [4] *Assume the underlying quiver contains no oriented cycles (and k is a field of characteristic zero). Then Γ has a unique source.*

The following counter-example shows that this result is not true.


Recall from [2] that given a bypass (α, u) in a quiver Q , then $\phi_{\alpha, u, \tau}$ denotes the automorphism of kQ which sends α to $\alpha + \tau u$, $\tau \in k \setminus \{0\}$, and leaves the other arrows fixed.

Key words and phrases. Fundamental group, bound quiver, triangular algebra.

Counter-example: Let k be a field of characteristic not equal 2, and $A = kQ/I$, where Q is the quiver $3 \begin{smallmatrix} \xrightarrow{\beta_1} \\ \xleftarrow{\beta_1} \end{smallmatrix} 2 \begin{smallmatrix} \xrightarrow{\alpha_1} \\ \xleftarrow{\alpha_2} \end{smallmatrix} 1$, and $I = \langle \alpha_1\beta_1, \alpha_2\beta_2 \rangle$. Since I is monomial, the fundamental group $\pi_1(Q, I)$ is isomorphic to the free group in two generators, $\mathbb{Z} \amalg \mathbb{Z}$. A straightforward computation shows that $kQ/I \simeq kQ/I_1 \simeq kQ/I_2 \simeq kQ/I_3 \simeq kQ/I_4$, where:

- $I_1 = \langle (\alpha_1 - \alpha_2)\beta_1, \alpha_2\beta_2 \rangle = \phi_{\alpha_1, \alpha_2, -1}(I)$, and leads to a fundamental group isomorphic to $\mathbb{Z}$;
- $I_2 = \langle \alpha_1(\beta_1 - \beta_2), \alpha_2\beta_2 \rangle = \phi_{\beta_1, \beta_2, -1}(I)$, and leads to a fundamental group isomorphic to $\mathbb{Z}$;
- $I_3 = \langle \alpha_1\beta_1 - \alpha_2\beta_1 - \alpha_1\beta_2, \alpha_2\beta_2 \rangle = \phi_{\alpha_1, \alpha_2, -1}(I_2) = \phi_{\beta_1, \beta_2, -1}(I_1)$, and leads to a trivial fundamental group;
- $I_4 = \langle \alpha_1\beta_1 + \alpha_2\beta_2, \alpha_2\beta_1 + \alpha_1\beta_2 \rangle$. The ideal I_4 leads to a fundamental group isomorphic to $\mathbb{Z}_2$ and is obtained from the automorphism ϕ of kQ defined by $\phi(\alpha_1) = \frac{1}{2}(\alpha_1 - \alpha_2)$, $\phi(\alpha_2) = \frac{1}{2}(\alpha_1 + \alpha_2)$, $\phi(\beta_1) = \frac{1}{2}(\beta_1 - \beta_2)$, $\phi(\beta_2) = (\beta_1 + \beta_2)$.

Moreover, the associated quiver Γ is then



Of course, there is a surjective group homomorphism $f : \pi_1(Q, I) \rightarrow \pi_1(Q, I_4)$. However, there is no path from $\sim_I$ to $\sim_{I_4}$ in Γ .

References

- [1] P. Le Meur. The fundamental group of an algebra without double bypasses. *C. R. Acad. Sci. Paris. Sér I*, 341, 2005.
- [2] P. Le Meur. The universal cover of an algebra without double bypass. *J. Algebra*, 312(1):330–353, 2007.
- [3] P. Le Meur. The universal cover of a monomial triangular algebra without multiple arrows. *J. Algebra Appl.*, 7(4):443–469, 2008.
- [4] F. Xu. A note on the fundamental group of a triangular algebra. *Algebra Coll*, 15(3):371–378, 2008.

J. C. BUSTAMANTE; DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD SAN FRANCISCO DE QUITO, QUITO, ECUADOR.

E-mail address: `juanb@usfq.edu.ec`

D. CASTONGUAY; INSTITUTO DE INFORMÁTICA, UNIVERSIDADE FEDERAL DE GOIÁS, GOIÂNIA, BRASIL.

E-mail address: `diane@inf.ufg`